

5-23-2016

Erosion of Empathy in Primary Care Trainees

Jacquelyn McRae, PharmD

Jefferson College of Population Health, Thomas Jefferson University, Jacquelyn.McRae@jefferson.edu

G. Calusi

AUSL

Mengdan Liu

Center for Research in Medical Education and Health Care, Thomas Jefferson University, mengdan.liu@jefferson.edu

V. Scognamiglio

AUSL

E. Messina

AUSL

*See next page for additional authors*Follow this and additional works at: <http://jdc.jefferson.edu/jcphposters> Part of the [Health Services Research Commons](#)

Recommended Citation

McRae, PharmD, Jacquelyn; Calusi, G.; Liu, Mengdan; Scognamiglio, V.; Messina, E.; Polenzani, L.; and Maio, PharmD, MSPH, Vittorio, "Erosion of Empathy in Primary Care Trainees" (2016). *Jefferson College of Population Health Posters*. 6.
<http://jdc.jefferson.edu/jcphposters/6>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Jefferson College of Population Health Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Jacquelyn McRae, PharmD; G. Calusi; Mengdan Liu; V. Scognamiglio; E. Messina; L. Polenzani; and Vittorio Maio, PharmD, MSPH



EROSION OF EMPATHY IN PRIMARY CARE TRAINEES

McRae J¹, Calusi G², Liu M³, Scognamiglio V², Messina E², Polenzani L², Maio V¹

1. Jefferson College of Population Health, Thomas Jefferson University, Philadelphia, PA 19107 United States

2. AUSL 4 Prato, Prato, Italy

3. Center for Medical Research and Education, Thomas Jefferson University, Philadelphia, PA 19107 United States

INTRODUCTION

- Empathy outside of the clinical context is oftentimes described as “feelings” or emotionally putting oneself in another’s situation (Halpern, 2014).
- Empathy in patient care is “a predominantly cognitive (as opposed to affective or emotional) attribute that involves an understanding (as opposed to feeling) of patients’ experiences, concerns, and perspectives combined with a capacity to communicate this understanding”(Hojat, 2009).
- Evidence supports that empathic care is associated with better health outcomes, enhanced shared-decision making, and higher rates of medication adherence, and lower rates of medical malpractice (Hojat, 2011; Hickson, 2002; Riess, 2012, Del Canale, 2012).
- Despite the positive effects of empathy, physicians struggle with exactly how to display empathy in the patient-provider relationship (Gleichgerrcht and Decety, 2012).
- To this end, educating physicians on how to demonstrate empathy in the early stages of their education may be an effective strategy to increasing empathic engagement.
- The evidence shows that, worrisomely, empathy among medical students tends to decrease with increased exposure to the healthcare system (Hojat, 2009).
- Compared to studies evaluating empathy in medical students, fewer studies have evaluated empathy in physician residents (Avasarala, 2015; Greenberg, 2015; Olson, 2015).
- Current evidence is conflicting as to whether empathy decreases or is unaffected during residency training (Greenberg, 2015; West, 2007).

OBJECTIVE

To evaluate if empathy among physician residents (trainees) differs dependent on training year and to assess trainees’ characteristics associated with higher empathy scores.

METHODS

This study was reviewed by the Thomas Jefferson University IRB and determined not to constitute human subjects research.

Setting

- In order to become a primary care physician (PCP) and practice for the Italian National Health Service, a physician must complete a three-year postgraduate training in primary care structured and managed by the 21 Italian regions.
- The training entails 4,800 hours of professional training of which one third are devoted to didactics and two thirds are dedicated to practical experience, including clinical activities in hospital and outpatient settings, with a focus on PCP ambulatory practice.
- In Tuscany, every year approximately 78 physicians enter the three-year regional postgraduate primary care training; the training is organized by the four regional local health authorities.

Instrument

The adapted Jefferson Scale of Physician Empathy (JSE) includes information on two domains:

- **Demographic information:** age, gender, years of training, and years of service for trainees.
- **Empathic measurement:** using the Italian translated, HP-version of JSE. A prior study confirmed the psychometrics of the Italian translation of the JSE (Di Lillo, 2009). The HP-version of JSE consists of twenty Likert questions on a seven point scale (1=strongly disagree, 7= strongly agree), with ten questions reverse coded. The total empathy score may range from a minimum of 20 to a maximum of 140.

Data collection

- From June to July, 2015, all 119 PCP trainees attending the PCP training program managed by the Florence and of Prato LHAs, were invited to complete an online version of the adapted JSE.
- A total of 98 trainees returned the survey (response rate: 82.4%).

Data analysis

- Descriptive statistics were generated for the demographic variables (Table 1).
- Mean empathy scores and standard deviation (SD) were calculated for trainees according to the demographic variables of interest (Table 2).
- Student’s t-test and analysis of variance (ANOVA) were conducted to evaluate statistical differences in mean empathy scores by subcategories, as appropriate. We used the Tukey’s Studentized Range Tests for statistically significant ANOVA results (Table 2).
- All statistical analyses were completed using SAS 9.3 Enterprise Software.

RESULTS

Table 1. Demographics for Sample Population (N=98)

	n (%), 2015
Age, yrs ^a	
Range	26-39
Mean (SD)	29 (2.7)
Median	29
Sex, n (%) ^a	
Female	66 (69%)
Male	30 (31%)
Trainee year, n (%) ^b	
First	35 (36%)
Second	28 (29%)
Third	35 (36%)
Years of service, n (%) ^a	
≤2	43 (45%)
3-5	46 (48%)
≥6	7 (7%)

Abbreviations: SD, standard deviation
a Available for 96 respondents
b Sum is not 100 due to rounding

Table 2. JSE Scores (N = 98)

Characteristic	n	Mean score (SD)	Range of scores	P value
All respondents	98	112.7 (12.7)	77-140	
Age group, yrs ^a				
26-28	36	116.8 (11.3)	87-133	0.036
29-31	40	109.4 (14.4)	77-140	
>32	20	111.5 (10.0)	97-133	
Sex ^a				
Female	66	113.5 (11.9)	77-140	0.324
Male	30	110.7 (14.6)	85-133	
Trainee year ^a				
First	35	116.0 (11.3)	87-132	0.035
Second	28	114.0 (13.1)	83-140	
Third	35	108.4 (12.8)	77-129	
Years of service ^a				
≤2	43	115.2 (11.5)	87-133	0.182
3-5	46	110.3 (14.3)	77-140	
≥6	7	111.6 (6.0)	102-120	

Abbreviations: ANOVA, analysis of variance; SD, standard deviation
a ANOVA P value for 3-way comparisons and t test P value for 2-way comparisons
Bolded values are statistically significant at alpha level of 0.05.

LIMITATIONS

- Our sample included physicians in a single primary care training program therefore, results may not be generalizable to primary care trainees across Italy or other geographic regions.
- Our results are from a cross-sectional study and therefore we did not capture the true variation of empathy as trainees progressed in the program.

DISCUSSION

- To our knowledge, this study is the first to investigate characteristics linked to empathic care in Italian primary care trainees and adds to the scarce knowledge on this topic.
- Our study showed empathy scores were significantly lower in year 3 of residency training. These results are in agreement with previous evidence that demonstrated that empathy within medical residents significantly declined during matriculation in training programs (Bellini, 2001; Bellini, 2005; West, 2006; West, 2007; Rosen, 2006; Mangione, 2002).
- These results are important because they demonstrate that primary care residents may follow the same phenomenon as medical students and internal medicine residents of decreasing empathy with increased exposure to clinical practice.
- We observed higher empathy scores in women vs. men, although not statistically significant likely due to small sample size (Park, 2015; Hojat, 2002).
- Future studies in larger samples may confirm our results. In addition, future research should utilize a longitudinal analysis of changes in empathy among primary care trainees.
- We urge directors of primary care residency programs to support educational activities to teach trainees strategies for empathic engagement.

REFERENCES

1. Halpern J. From idealized clinical empathy to empathic communication in medical care. Med Health Care and Philos. 2014; 12: 301-311.
2. Hojat M, Gonnella JS, Nasca TJ, Mangione S, Vergare M, and Magee M. Physician Empathy: Definition, Components, Measurement, and Relationship to Gender and Specialty. Am J Psychiatry. 2002; 159: 1563-1569.
3. Aring, C. Sympathy and Empathy. JAMA. 1958; 167(4):448-452.
4. Hojat M, Vergare MJ, Maxwell GB, Jerrine S, Isenberg GA, Veloski J, and Gonnella JS. The Devils is in the Third Year: A Longitudinal Study of Erosion of Empathy in Medical School. Acad Med. 2009; 84:1182-91.
5. Hojat M, Louis DZ, Markham FW, Wender R, Rabinowitz C, and Gonnella JS. 2011. Physicians' empathy and clinical outcomes for diabetic patients. Academic Medicine 86(3): 359.
6. Hickson GB, Federspiel CF, Pichert JW, Miller CS, Gauld-Jaeger J, Bost P. Patient complaints and malpractice risk. JAMA. 2002; 287(22): 2951–7. doi: 10.1001/jama.287.22.2951.
7. Riess H, Kelley JM, Bailey RW, Dunn EJ, Phillips M. Empathy training for resident physicians: a randomized controlled trial of a neuroscience-informed curriculum. J Gen Intern Med. 2012;27(10): 1280–6. doi: 10.1007/s11606-012-2063-z
8. Del Canale S, Louis DZ, Maio V, Wang X, Rossi G, Hojat M, Gonnella JS. Physicians' Empathy and Disease Complications: An Empirical Study of Primary Care Physicians and Their Diabetic Patients in Parma, Italy Academic Medicine 2012;87(9), 1243-1249).
9. Gleichgerrcht E, Decety J (2012) The costs of empathy among health professionals. In: Decety J, editor. Empathy: from Bench to Bedside. Cambridge: MIT Press. 245–2611.
10. Avasarala SK, Whitehouse S, Drake SM. Internship and Empathy: Variations Across Time and Specialities. South Med J. 2015; 108(10):591-5. doi:10.14423/SMJ.0000000000000347.
11. Greenberg L., Agrawal D., Toto R., Blatt B. Empathy, Sense of Power, and Personality Do They Change During Pediatric Residency? South Med J. 2015; 108(8):471-4.
12. Olson K, Kemper KJ, Mahan JD. What factors promote resilience and protect against burnout in first-year pediatric and medicine-pediatric residents? J Evid Based Complementary Altern Med. 2015; 20(3):192-8. doi: 10.1177/2156587214568894.